

A single-base substitution within an intronic repetitive element causes dominant retinitis pigmentosa with reduced penetrance

Thomas Rio Frio¹, Terri L. McGee², Nicholas M. Wade¹, Christian Iseli³, Jacques S. Beckmann^{1,4}, Eliot L. Berson², Carlo Rivolta^{1*}

¹Department of Medical Genetics, University of Lausanne, Lausanne, Switzerland

²The Berman-Gund Laboratory for the Study of Retinal Degenerations, Harvard Medical School, Massachusetts Eye and Ear Infirmary, Boston, Massachusetts, USA

³Ludwig Institute for Cancer Research and Swiss Institute of Bioinformatics, Lausanne, Switzerland

⁴Service of Medical Genetics, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland

email: Carlo Rivolta (carlo.rivolta@unil.ch)

*Correspondence to Carlo Rivolta, Department of Medical Genetics, University of Lausanne, Rue du Bugnon 27, 1005 Lausanne, Switzerland

Communicated by Andreas Gal

Keywords

PRPF31 • RP11 • splicing • UHT sequencing

Abstract

We report the study of a large American family displaying autosomal dominant retinitis pigmentosa with reduced penetrance, a form of hereditary retinal degeneration. Although the inheritance pattern and previous linkage mapping pointed to the involvement of the PRPF31 gene, extensive screening of all its exons and their boundaries failed in the past to reveal any mutation. In this work, we sequenced the entire PRPF31 genomic region by both the classical Sanger method and ultrahigh throughput (UHT) sequencing. Among the many variants identified, a single-base substitution (c.1374+654C>G) located deep within intron 13 and inside a repetitive DNA element was common to all patients and obligate asymptomatic carriers. This change created a new splice donor site leading to the synthesis of two mutant PRPF31 isoforms, degraded by nonsense-mediated mRNA decay. As a consequence, amounts of PRPF31 mRNA derived from the mutant allele were very reduced, with no evidence of mutant proteins being synthesized. Our results indicate that c.1374+654C>G causes retinitis pigmentosa via haploinsufficiency, similar to the vast majority of PRPF31 mutations described so far. We discuss the potential of UHT sequencing technologies in mutation screening and the continued identification of pathogenic splicing mutations buried deep within intronic regions.

Hum Mutat 30:1-8, 2009.

© 2009 Wiley-Liss, Inc.

Received: 19 January 2009; Accepted: 8 June 2009

Digital Object Identifier (DOI) 10.1002/humu.21071